

Structure of the Electric Flux in $\mathcal{N} = 4$ Supersymmetric Yang-Mills Theory

J. Erickson*, G.W. Semenoff* and K. Zarembo*,[†]

*The Niels Bohr Institute
Blegdamsvej 17
DK-2100 Copenhagen Ø
Denmark*

Abstract

Correlators of Wilson loop operators with $\mathcal{O}_4 = \text{Tr}(F_{\mu\nu}^2 + \dots)$ are computed in $\mathcal{N} = 4$ super-Yang-Mills theory using the AdS/CFT correspondence. The results are compared with the leading order perturbative computations. As a consequence of conformal invariance, these correlators have identical forms in the weak and strong coupling limits for circular loops. They are essentially different for contours not protected by conformal symmetry.

* Department of Physics and Astronomy, University of British Columbia, 6224 Agricultural Road, Vancouver, British Columbia, Canada V6T 1Z1.

[†] Institute of Theoretical and Experimental Physics, B. Cheremushkinskaya 25, 117259 Moscow, Russia.

E-mail: erickson@nbi.dk, semenoff@nbi.dk, zaremba@itep.ru

The conjectured duality of $\mathcal{N} = 4$, $D = 4$ super-Yang-Mills theory (SYM) to supergravity on $AdS_5 \times S_5$ gives rise to the possibility of obtaining quantitative information about the former theory in the limit of infinite 't Hooft coupling [1, 2, 3, 4]. Though this is a regime which is inaccessible to conventional perturbative field theory methods, some of the results obtained from the AdS/CFT correspondence can be tested by non-renormalization theorems which protect certain correlation functions from gaining radiative corrections [4][‡]. The quantities not protected by supersymmetry probe the genuine strong-coupling dynamics of the SYM theory. An example of such a quantity is the interaction potential between static charges. The Coulomb form of the potential is dictated by the conformal invariance of $\mathcal{N} = 4$ SYM, but the strength of the interaction is not fixed by any symmetry and appears to be proportional to $\sqrt{g^2 N}$ at strong coupling [16, 17] in contrast to the weak-coupling $O(g^2 N)$ behavior.

The purpose of the present paper is to study the structure of the electric field induced by static charges in the vacuum. The static charges are introduced by a Wilson loop. The one appropriate for supergravity calculations corresponds to the phase factor acquired by an infinitely heavy W-boson propagating in background gauge and scalar fields [16, 18]:

$$W(C) = \frac{1}{N} \text{Tr P exp} \oint_C d\tau (iA_\mu \dot{x}_\mu + \Phi_i \dot{y}_i), \quad (1)$$

where the A_μ are $U(N)$ gauge fields and the Φ_i are scalars in the adjoint representation, both represented by Hermitian matrices. Moreover, $\dot{y}_i = \sqrt{\dot{x}^2} \theta_i$ where θ_i is a point on the five-dimensional sphere: $\theta^2 = 1$.

The form of the coupling to scalars in (1) is natural from several points of view [18, 19]. In particular, this Wilson loop operator is annihilated by half of the supercharges if C is a straight line [18]. On the other hand, closed Wilson loops are generally not supersymmetric. A circular loop, however, is a special case since it can be mapped to a straight line by a conformal transformation. The straight line Wilson loop operator commutes with half of the supersymmetry generators. Thus, circular loops are annihilated by a combination of supercharges twisted by conformal transformations[§]:

$$\tilde{Q}_\alpha = \Omega(C)^\dagger Q_\alpha \Omega(C), \quad (2)$$

where $\Omega(C)$ is a unitary operator implementing a conformal transformation that maps C to a line. Since conformal invariance is an exact symmetry of the $\mathcal{N} = 4$ SYM vacuum, $\Omega|0\rangle = |0\rangle$ and the twisted supercharges annihilate the vacuum: $\tilde{Q}_\alpha|0\rangle = 0$, along with the original supersymmetry generators Q_α . If the Wilson loop has a different geometry,

[‡]Various non-renormalization theorems for $\mathcal{N} = 4$ SYM are discussed in [5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15].

[§]In discussing supersymmetry, it is convenient to consider $\mathcal{N} = 4$ SYM theory as a reduction of ten-dimensional $\mathcal{N} = 1$ theory. Then the supercharges are the components of a ten-dimensional Majorana-Weyl spinor Q_α . The gauge potentials and the scalar fields form a ten-dimensional vector: $A_M = (A_\mu, \Phi_i)$. The supercharges act on gauge potentials as $[Q_\alpha, A_M] = \frac{i}{2}(\Gamma_M \Psi)_\alpha$, where Γ_M are Dirac matrices and Ψ is the ten-dimensional Majorana-Weyl fermion, the reduction of which to four dimensions gives the fermionic content of $\mathcal{N} = 4$ SYM theory. The supersymmetry transformation of a Wilson loop produces degenerate combination of the Dirac matrices [18]: $\bar{\epsilon} \dot{X}^M \Gamma_M \Psi$, $X^M = (x^\mu, iy^i)$, since $\dot{X}^2 = 0$. An appropriate choice of ϵ turns the variation of a Wilson loop to zero if $\dot{X}^M$ is constant, that is, if C is a straight line.

generically there is no residual supersymmetry. This is the case for a loop consisting of two infinite anti-parallel lines; this loop corresponds to a static electric dipole.

The structure of electric field induced by the charges can be probed by local gauge invariant operators, like $\text{Tr } F_{\mu\nu}^2$. On the supergravity side,

$$\mathcal{O}_4(x) = \frac{1}{2g^2} \text{Tr} \left\{ F_{\mu\nu}^2 + [\Phi_i, \Phi_j]^2 + \bar{\psi} \Gamma^i [\Phi_i, \psi] \right\} \quad (3)$$

couples to the dilaton. We consider the correlator of the Wilson loop with $\mathcal{O}_4(x)$:

$$\mathcal{W}(C, x) = - \frac{\langle W(C) \frac{1}{2g^2} \text{Tr}(F_{\mu\nu}^2(x) + \dots) \rangle_{\text{conn}}}{\langle W(C) \rangle}, \quad (4)$$

where the dots denote the scalar and the fermionic terms required to put the operator in the short multiplet of $\mathcal{N} = 4$ SUSY. The correlator (4) probes the local electric field induced by non-dynamical charges whose trajectories form the loop C . This correlator has been discussed in the context of the AdS/CFT correspondence in the limiting case when the distance from the point x to the loop C is large and the Wilson loop can be expanded in local operators [20]. The correlator (4) was also used to demonstrate that, in the finite-temperature, confining phase of strongly coupled $\mathcal{N} = 4$ SYM, the classical string in the bulk is projected onto an electric flux tube, stretched between static charges, which has thickness on the order of the correlation length [21][¶]. In the conformal theory, the correlator must fall-off as a power of the distance from the loop. At weak coupling, the correlator decreases with distance as $1/d^6$, since the classical electric field of the dipole falls off as $1/d^3$. We shall show that the correlator obeys a different power law at strong coupling.

At strong coupling, correlators in the SYM theory are described by classical supergravity on $AdS_5 \times S_5$. We choose units in which the radius of AdS_5 is unity. The metric on $AdS_5 \times S_5$ in these units is

$$ds^2 = \frac{1}{z^2} (dz^2 + dx_\mu dx_\mu) + d\Omega_5^2. \quad (5)$$

The Wilson loop on the boundary of AdS_5 at $z = 0$ creates the classical string in the bulk; it is the surface of minimal area bounded by the loop [16, 17]. The correlation function (4) is given by the dilaton propagator with one end attached to the point x and the other integrated over the minimal surface spanning the loop C [20]:

$$\mathcal{W}_{g \rightarrow \infty} \propto \sqrt{g^2 N} \int d^2 \sigma \sqrt{\det G} \frac{Z^4}{[Z^2 + |\vec{X} - \vec{x}|^2]^4}, \quad (6)$$

where $G_{ab} = g_{MN} \partial_a Y^M \partial_b Y^N$ is the metric induced on the minimal surface. The surface is parameterized by coordinates $Y^M = (Z(\sigma), X^\mu(\sigma))$ in AdS_5 .

At weak coupling, the correlator (4) is given by the lowest-order Feynman diagram:

$$\begin{aligned} \mathcal{W}_{g \rightarrow 0} = & \frac{g^2 N}{4} \oint_C dy_\mu \oint_C dz_\nu \left[\delta_{\mu\nu} \frac{\partial}{\partial x_\lambda} \frac{1}{4\pi^2 |\vec{x} - \vec{y}|^2} \frac{\partial}{\partial x_\lambda} \frac{1}{4\pi^2 |\vec{x} - \vec{z}|^2} \right. \\ & \left. - \frac{\partial}{\partial x_\nu} \frac{1}{4\pi^2 |\vec{x} - \vec{y}|^2} \frac{\partial}{\partial x_\mu} \frac{1}{4\pi^2 |\vec{x} - \vec{z}|^2} \right]. \end{aligned} \quad (7)$$

[¶]More applications are discussed in [22].

We start with the supersymmetry preserving circular loop. At weak coupling, we find:

$$\mathcal{W}_{g \rightarrow 0} = \frac{g^2 N}{4\pi^2} \frac{R^4}{[(y^2 + r^2 - R^2)^2 + 4R^2 y^2]^2}, \quad (8)$$

where R is the radius of the loop, r is the radial coordinate of the point x , and y is the distance from x to the plane of the loop. This expression coincides with the result of the strong-coupling supergravity calculation given in [20]:

$$\mathcal{W}_{g \rightarrow \infty} \propto \sqrt{g^2 N} \frac{R^4}{[(y^2 + r^2 - R^2)^2 + 4R^2 y^2]^2}. \quad (9)$$

The functional form of the correlator of $\text{Tr}(F^2 + \dots)$ with the circular loop appears not to be renormalized. Actually, this non-renormalization property is a consequence of conformal symmetry. Although the scale invariance only fixes the above correlator up to a function of two dimensionless ratios of the parameters r , R and y , the conformal transformation which maps the circle to the line reduces the number parameters to one, the distance d between the point and the line. The correlator of $\text{Tr}(F_{\mu\nu}^2 + \dots)$ with the Wilson line is determined by the dimension of the dilaton operator and is proportional $1/d^4$. Transforming the line to the circle we get (8) with an overall coefficient which can depend only on the coupling. The distinguished role of the supersymmetry preserving circular loops is reflected in the fact that minimal surface in AdS_5 bounded by it has a very simple form [18, 20]. The minimal surfaces bounded by contours which do not have such nice conformal properties are generically more complicated [16, 17, 23, 18]. We will find that for such contours the electric field profile probed by the correlator (4) is renormalized in a non-trivial way.

We consider the loop composed of antiparallel infinite lines. The leading-order perturbative result for the correlator with the dilaton operator in this case is

$$\begin{aligned} \mathcal{W}_{g \rightarrow 0} &= \frac{g^2 N}{64\pi^2} \left\{ \frac{1}{[y^2 + (d - L/2)^2]^2} - 2 \frac{y^2 + d^2 - L^2/4}{[y^2 + (d - L/2)^2]^{3/2} [y^2 + (d + L/2)^2]^{3/2}} \right. \\ &\quad \left. + \frac{1}{[y^2 + (d + L/2)^2]^2} \right\} \\ &= \frac{g^2 N}{4} \left[\frac{\partial}{\partial x_i} \left(\frac{1}{4\pi|\mathbf{x} - \mathbf{L}/2|} - \frac{1}{4\pi|\mathbf{x} + \mathbf{L}/2|} \right) \right]^2, \end{aligned} \quad (10)$$

where L is the distance between the lines; d is the distance, in the plane of the lines, between the point x and the line midway between the two lines; and y is the distance between x and the plane of the lines: $\mathbf{L} = (0, L, 0, 0)$, $\mathbf{x} = (0, d, y, 0)$.

An expression for the minimal surface whose boundary is the two lines is given in [16]:

$$x = Z_0 \int_1^{Z_0/Z} \frac{du}{u^2 \sqrt{u^4 - 1}}, \quad (11)$$

with

$$Z_0 = \frac{L}{2} \left[\int_1^\infty \frac{du}{u^2 \sqrt{u^4 - 1}} \right]^{-1} = \frac{L}{2} \frac{\Gamma(1/4)^2}{\sqrt{2}\pi^{3/2}}. \quad (12)$$




Figure 1: $\mathcal{W}_{g \rightarrow \infty}$ (the shorter peak) and $\mathcal{W}_{g \rightarrow 0}$ (the taller peak), drawn as a function of y with $L = 1$ and $d = 0$. The heights are normalized so that the area under each curve is equal to unity. Note that asymptotically the $\mathcal{W}_{g \rightarrow \infty}$ falloff is actually faster (y^{-7} versus y^{-6}).

From (6) we find:

$$\begin{aligned} \mathcal{W}_{g \rightarrow \infty} &\propto \sqrt{g^2 N} Z_0^2 \int_{-\infty}^{\infty} dt \int_{-L/2}^{L/2} dx \frac{1}{[Z^2 + y^2 + (x-d)^2 + t^2]^4} \\ &\propto \sqrt{g^2 N} L^2 \int_{-L/2}^{L/2} dx \frac{1}{[Z^2 + y^2 + (x-d)^2]^{7/2}}, \end{aligned} \quad (13)$$

where y , L , and d are as before.

This result is clearly different than that of the perturbative calculation. In particular, we can consider the asymptotic behavior of each result when the distance to the point x is much larger than the separation between the charges L . Expanding (13) in L gives

$$\mathcal{W}_{g \rightarrow \infty} \propto \frac{\sqrt{g^2 N} L^3}{(y^2 + d^2)^{7/2}} + O(L^5). \quad (14)$$

The perturbative correlator (10) is just the field strength of electric dipole squared and is proportional to L^2 for small L :

$$\mathcal{W}_{g \rightarrow 0} = \frac{g^2 N}{64\pi^2} \frac{(4d^2 + y^2)L^2}{(y^2 + d^2)^4} + O(L^4). \quad (15)$$

Thus, the two correlators obey different power laws at large d , with $\mathcal{W}_{g \rightarrow \infty}$ falling off as d^{-7} and $\mathcal{W}_{g \rightarrow 0}$ falling off as d^{-6} . Figure 1 shows the form of the correlator (4), as a function of y , at strong and weak coupling, for $d = 0$.

However, the leading singularity as the point x approaches the loop obeys the same power law in the weak and the strong coupling limits. In both cases, the correlator behaves as ϵ^{-4} , where $\epsilon = d - L/2$ is the distance from the loop, as can be checked from (10) and (13). This is to be expected as the dimension of the dilaton operator is not renormalized.

To conclude, we have calculated correlators of Wilson loops with the dimension four chiral operator $\text{Tr}(F^2 + \dots)$ in $\mathcal{N} = 4$ SYM. We have compared the perturbative expression with

the strong coupling result predicted by the AdS/CFT correspondence. The correlator for a circular loop has the same functional form in both regimes, whereas the answer is completely different for antiparallel Wilson lines. Even the asymptotic form of the correlator at infinity appears to be modified.

Finally, we observe that the coupling constant dependence of $\mathcal{W}(C, x)$ at large N is natural and is related to the strong coupling limit of the Wilson loop expectation value. Indeed, $\mathcal{O}_4$ is related to the action density of super-Yang-Mills theory if we add terms which are a total divergence, and use the equations of motion for the scalar and fermion fields. Thus, the integral of (4) over the coordinate point x is related to the derivative of the expectation value of the Wilson loop by the coupling constant,

$$\int d^4x \mathcal{W}(C, x) = g^2 \frac{\partial}{\partial g^2} \ln W(C) \quad (16)$$

Also, we see from (6) that, in the $g \rightarrow \infty$ limit, this integral of $\mathcal{W}(C, x)$ is equal to $\pi^2/6$ times the minimal area of the surface. Thus, if the Wilson loop correlator at strong coupling is $\ln W(C) = \sqrt{4\pi g_{YM}^2 N}$ then we would expect that $\mathcal{W}(C, x) = 3\sqrt{4\pi g_{YM}^2 N}/\pi^2$.

Acknowledgments

We are grateful to Y. Makeenko and M. van Raamsdonk for discussions and to J. Ambjørn for hospitality at the Niels Bohr Institute. The work of KZ was supported by NATO Science Fellowship and, in part, by INTAS grant 96-0524, RFFI grant 97-02-17927 and grant 96-15-96455 for the promotion of scientific schools. GWS was supported by NSERC of Canada, the Niels Bohr Fund of Denmark and NATO CRG 970561.

References

- [1] J. Maldacena, “The Large N limit of superconformal field theories and supergravity,” Adv. Theor. Math. Phys. **2**, 231 (1998) hep-th/9711200.
- [2] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, “Gauge theory correlators from non-critical string theory,” Phys. Lett. **B428**, 105 (1998) hep-th/9802109.
- [3] E. Witten, “Anti-de Sitter space and holography,” Adv. Theor. Math. Phys. **2**, 253 (1998) hep-th/9802150.
- [4] O. Aharony, S.S. Gubser, J. Maldacena, H. Ooguri and Y. Oz, “Large N field theories, string theory and gravity,” hep-th/9905111.
- [5] S.S. Gubser and I.R. Klebanov, “Absorption by branes and Schwinger terms in the world volume theory,” Phys. Lett. **B413**, 41 (1997) hep-th/9708005.
- [6] D. Anselmi, D.Z. Freedman, M.T. Grisaru and A.A. Johansen, “Nonperturbative formulas for central functions of supersymmetric gauge theories,” Nucl. Phys. **B526**, 543 (1998) hep-th/9708042.

- [7] D.Z. Freedman, S.D. Mathur, A. Matusis and L. Rastelli, “Correlation functions in the CFT(d) / AdS(d+1) correspondence,” Nucl. Phys. **B546**, 96 (1999) hep-th/9804058.
- [8] S. Lee, S. Minwalla, M. Rangamani and N. Seiberg, “Three point functions of chiral operators in $D = 4$, $N=4$ SYM at large N ,” Adv. Theor. Math. Phys. **2**, 697 (1998) hep-th/9806074.
- [9] E. D’Hoker, D.Z. Freedman and W. Skiba, “Field theory tests for correlators in the AdS / CFT correspondence,” Phys. Rev. **D59**, 045008 (1999) hep-th/9807098.
- [10] K. Intriligator, “Bonus symmetries of $N=4$ superYang-Mills correlation functions via AdS duality,” hep-th/9811047..
- [11] F. Gonzalez-Rey, B. Kulik and I.Y. Park, “Nonrenormalization of two point and three point correlators of $N=4$ SYM in $N=1$ superspace,” hep-th/9903094.
- [12] K. Intriligator and W. Skiba, “Bonus symmetry and the operator product expansion of $N=4$ SuperYang-Mills,” hep-th/9905020.
- [13] B. Eden, P.S. Howe and P.C. West, “Nilpotent invariants in $N=4$ SYM,” hep-th/9905085.
- [14] A. Petkou and K. Skenderis, “A Nonrenormalization theorem for conformal anomalies,” hep-th/9906030.
- [15] H. Liu and A.A. Tseytlin, “Dilaton - fixed scalar correlators and $AdS_5 \times S^5$ - SYM correspondence,” hep-th/9906151.
- [16] J. Maldacena, “Wilson loops in large N field theories,” Phys. Rev. Lett. **80**, 4859 (1998) hep-th/9803002.
- [17] S. Rey and J. Yee, “Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity,” hep-th/9803001.
- [18] N. Drukker, D.J. Gross and H. Ooguri, “Wilson loops and minimal surfaces,” hep-th/9904191.
- [19] Y.M. Makeenko, “Polygon Discretization Of The Loop Space Equation,” Phys. Lett. **212B**, 221 (1988).
- [20] D. Berenstein, R. Corrado, W. Fischler and J. Maldacena, “The Operator product expansion for Wilson loops and surfaces in the large N limit,” Phys. Rev. **D59**, 105023 (1999) hep-th/9809188.
- [21] U.H. Danielsson, E. Keski-Vakkuri and M. Kruczenski, “Vacua, propagators, and holographic probes in AdS / CFT,” JHEP **01**, 002 (1999) hep-th/9812007.
- [22] C.G. Callan and A. Güijosa, “Undulating strings and gauge theory waves,” hep-th/9906153.
- [23] K. Zarembo, “Wilson loop correlator in the AdS / CFT correspondence,” hep-th/9904149.